



# Place of Residence at assessment, one month and three months



# Level of Care Changes



## Reasons for change

Had expected outcome scores, chose to stay at home at time of assessment, but later decided to move to residential care

Hospitalisation and decline in baseline function necessitating reassessment

Improvement in overall function and client decision to return home from residential care




Point of note...


**None of the clients indicated as needing rest home level care were referred back for increase to hospital level care within the three month period**



# Triggers



# Behavioural, Clinical and other issues indicated by MAPLe



# Exceptions

69yr old lady

Metastatic breast cancer

Not an exception by outcome scores however, support needs could have been met at home.

Except for home being beyond repair for her to be able to safely return home

ADL Hierarchy Scale (0-6)	2
Body Mass Index	0
Cognitive Performance Scale (0-6)	1
Communication Scale (0-8)	0
Depression Rating Scale (0-14)	0
IADL Capacity (0-48)	33
IADL Performance (0-48)	34
Pain Scale (0-4)	1
CHESS (0-5)	2
MAPLE Score (1-5)	4
Aggressive Behaviour Scale (0-12)	0
Pressure Ulcer Risk (0-8)	0
ADL Short Form (0-16)	2
ADL Long Form (0-28)	7



# Exceptions

90yr old lady with metastatic cancer

ADL Long form and CPS suggest rest home level care, decision made for hospital level care given the combination of cognition and ADL needs, quickly deteriorating medical condition.

Passed away within one month from assessment

ADL Hierarchy Scale (0-6)	2
Body Mass Index	0
Cognitive Performance Scale (0-6)	3
Communication Scale (0-8)	1
Depression Rating Scale (0-14)	0
IADL Capacity (0-48)	42
IADL Performance (0-48)	42
Pain Scale (0-4)	2
CHESS (0-5)	1
MAPLE Score (1-5)	5
Aggressive Behaviour Scale (0-12)	0
Pressure Ulcer Risk (0-8)	1
ADL Short Form (0-16)	6
ADL Long Form (0-28)	12





# Exceptions

65yr old gentleman with paranoid syndrome and persecutory phenomena secondary to Parkinsons and medications.

Persistent anxiety that he would harm his wife or himself

Captured in Depression Rating Scale

ADL Hierarchy Scale (0-6)	0
Body Mass Index	0
Cognitive Performance Scale (0-6)	1
Communication Scale (0-8)	1
Depression Rating Scale (0-14)	4
IADL Capacity (0-48)	27
IADL Performance (0-48)	27
Pain Scale (0-4)	0
CHESS (0-5)	2
MAPLE Score (1-5)	3
Aggressive Behaviour Scale (0-12)	0
Pressure Ulcer Risk (0-8)	1
ADL Short Form (0-16)	0
ADL Long Form (0-28)	0



# Exceptions

79yr old lady with cognitive decline, anxiety and alcohol abuse.  
Lack of insight into her own needs causing self neglect, but also neglect of her husband with whom she lived and required significant care.

ADL Hierarchy Scale (0-6)	3
Body Mass Index	20
Cognitive Performance Scale (0-6)	2
Communication Scale (0-8)	0
Depression Rating Scale (0-14)	0
IADL Capacity (0-48)	26
IADL Performance (0-48)	9
Pain Scale (0-4)	2
CHESS (0-5)	2
MAPLE Score (1-5)	4
Aggressive Behaviour Scale (0-12)	0
Pressure Ulcer Risk (0-8)	1
ADL Short Form (0-16)	4
ADL Long Form (0-28)	4



# Exceptions

72yr old lady post fractured neck of femur

Fluctuating level of assistance and at times requiring 2 persons.

Pain an issue

ADL Hierarchy Scale (0-6)	2
Body Mass Index	0
Cognitive Performance Scale (0-6)	1
Communication Scale (0-8)	1
Depression Rating Scale (0-14)	1
IADL Capacity (0-48)	40
IADL Performance (0-48)	40
Pain Scale (0-4)	3
CHESS (0-5)	1
MAPLE Score (1-5)	4
Aggressive Behaviour Scale (0-12)	0
Pressure Ulcer Risk (0-8)	2
ADL Short Form (0-16)	5
ADL Long Form (0-28)	12



# Exceptions

72yr old gentleman with metastatic cancer, rheumatoid arthritis, anxiety, and depression.

Cognitive and ADL outcomes scores suggest rest home level, however end stage disease and paranoia (indicated by CHESS and DRS) also taken into consideration and therefore decision made for hospital level care

ADL Hierarchy Scale (0-6)	3
Body Mass Index	0
Cognitive Performance Scale (0-6)	3
Communication Scale (0-8)	2
Depression Rating Scale (0-14)	2
IADL Capacity (0-48)	42
IADL Performance (0-48)	42
Pain Scale (0-4)	2
CHESS (0-5)	4
MAPLE Score (1-5)	5
Aggressive Behaviour Scale (0-12)	0
Pressure Ulcer Risk (0-8)	3
ADL Short Form (0-16)	6
ADL Long Form (0-28)	10



# Exceptions

86yr old lady with diagnosis of anxiety

Moved in with her daughter from out of the area who had been advised that she already met residential care criteria.

	Dec 14	Mar 14
ADL Hierarchy Scale (0-6)	0	0
Body Mass Index	0	22
Cognitive Performance Scale (0-6)	2	2
Communication Scale (0-8)	0	0
Depression Rating Scale (0-14)	0	2
IADL Capacity (0-48)	1	13
IADL Performance (0-48)	1	24
Pain Scale (0-4)	0	2
CHESS (0-5)	1	0
MAPLE Score (1-5)	4	4
Aggressive Behaviour Scale (0-12)	0	0
Pressure Ulcer Risk (0-8)	0	1
ADL Short Form (0-16)	0	0
ADL Long Form (0-28)	0	0



# Thoughts on strengthening the tool

Perhaps the MAPLe would be better utilised as the vulnerability and risk indicator as all encompassing – however rather than assign a MAPLe value to a particular level perhaps a score of 3+ for any level of care.

Behavioural considerations: Aggressive Behaviour Scale

Depressive/Anxiety considerations: Depression Rating Scale

Clinical considerations:

CHESS score

Pain Scale

Pressure Ulcer Risk



## Closing Remarks

